


PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61N 5/06	A1	(11) International Publication Number: WO 98/08568 (43) International Publication Date: 5 March 1998 (05.03.98)
(21) International Application Number: PCT/GB97/02322 (22) International Filing Date: 29 August 1997 (29.08.97) (30) Priority Data: 9618051.8 29 August 1996 (29.08.96) GB (71) Applicant (for all designated States except US): SLS BIO-PHILE LIMITED [GB/GB]; Units 1 & 2, Heol Rhosyn, Dafen Industrial Estate, Llanelli, Carmarthenshire SA14 8LX (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): CLEMENT, Robert, Marc [GB/GB]; 11 Plas Road, Pontardawe, Swansea SA8 3HD (GB). KIERNAN, Michael, Noel [GB/GB]; 89 Heol Ed-dwch, Seven Sisters, Neath SA10 9AW (GB). (74) Agent: AUSTIN, Hedley, William; Urquhart-Dykes & Lord, Alexandra House, Alexandra Road, Swansea SA1 5ED (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: WRINKLE REMOVAL  (57) Abstract <p>Wrinkles are comestically removed from a superficial area of mammalian skin tissue having an epidermal layer, a basal layer, and a dermal layer, by irradiating the dermal layer through the basal layer by means of visible or infrared radiation (such as a laser), the irradiation being selected to be absorbed by a chromophore in the dermal layer such that collagen present in the dermal layer is heated, while the basal layer remains intact so as to substantially inhibit contact of the dermal layer with ambient air.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TC	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Wrinkle Removal

The present invention relates to a method of removing wrinkles from a superficial area of mammalian skin tissue.

The application of laser technology in healthcare is well known, and the use of lasers in medical applications has been studied extensively since the early 1960's. In recent years an increasing interest has been shown in cosmetic applications. One such cosmetic application is skin resurfacing and wrinkle removal; in this field lasers can be used as an alternative to surgical facelifts. A CO₂ laser, which operates in the infra-red region of the spectrum, is used, the radiation being totally absorbed in a thin layer of surface tissue, primarily because of the wavelength's high rate of absorption in water.

The current approach used to achieve CO₂ laser based skin resurfacing and wrinkle removal involves a two stage process, as follows:

Stage 1 - Skin Resurfacing: The purpose of the first stage is to remove a thin layer of tissue from a superficial area of mammalian skin tissue. To achieve this, the skin is irradiated by the laser which, as mentioned above, deposits its energy in a thin layer at the surface (approx 30 μ m in thickness). The application of the laser radiation to the tissue is by computer controlled scanner designed to ensure that one pass of the radiation removes one "layer" of tissue only. Thus damage is limited to the epidermis only. This process quickly ablates the surface layer removing the surface skin and with it any skin blemishes. The process is straightforward and does not demand a great deal of operator skill. The surface skin immediately starts to replace itself and this allows the skin to return to normal in about 3 to 5 days.

Stage 2 - Wrinkle Removal: After the skin surface has been resurfaced, the operator will, typically, apply further "passes" of the laser, thereby removing further layers of mammalian skin tissue, each layer being of the order of 30 μ m thick. Typically, around three or four further passes will complete the wrinkle removal procedure. This treatment therefore produces a controlled second degree burn. The skin barrier to the outside world is removed and therefore, there is a danger of infection.

The required effect is two fold:

- (a) the laser induces denaturing of the collagen in the dermis, and the formation of cross links, which results in a tightening effect stretching the skin, reducing or removing the wrinkles (it is thought that the thermal threshold for this effect is a temperature of 70°C); and
- (b) the changes to the dermis induce the generation of new collagen which develops using the matrix created by the denatured collagen as a foundation.

The skin-resurfacing and wrinkle removal procedure outlined above is considered by many experts in the field as a significant improvement over previously used surgical methods. The procedure uses the laser's ability to deliver high energy density at the surface of tissue and hence ablate the surface tissue in a well controlled manner. Continuing to remove the tissue, layer by layer is designed to damage the collagen and hence induce wrinkle removal. This second stage of the procedure is primitive; the skin weeps, scabs form and redness of the skin appears for many weeks.

It is therefore the aim of the present invention to provide a method of removing wrinkles from a superficial area of mammalian skin tissue without causing secondary burns and other problems associated with traditional wrinkle removal.

If the target for the laser has an appropriate chromophore (a substance that absorbs a specific wavelength and transmits or scatters at other wavelengths) then the laser can be used to modify that target selectively within an inhomogeneous volume of tissue. Occasionally, the desired target does not have a suitable chromophore of its own but exists in close proximity to another material which has such a chromophore which can be selectively targeted. Such interaction is called secondary selective interaction.

According to the present invention there is provided a method of removing wrinkles from a superficial area of mammalian skin tissue having, in the order specified, an epidermal layer, a basal layer, and a dermal layer, which method comprises: irradiating said dermal layer through said basal layer by means of visible or infra-red radiation, said irradiation being selected to be absorbed by a chromophore in said dermal layer such that collagen present in said dermal layer is heated, while said basal layer remains intact so as to substantially inhibit contact of said dermal layer with ambient air.

The irradiation of the dermal layer in the method according to the invention is therefore such as to shrink the skin tissue without damage to the dermis (in other words, without causing second degree burns) because the barrier provided by the basal layer remains intact. The method according to the invention is non-invasive and can readily be performed by non-medical personnel.

The irradiation may be by means of a source of visible or infra-red radiation (suitably filtered to remove deleterious ultra-violet radiation if necessary). The radiation may be incoherent or, more preferably, coherent (that is from a laser source). Such a laser source may be, for example, a dye laser, a ruby laser, or a semi-conductor laser. If a dye laser is used, its wavelength is preferably such that it is absorbed by oxyhemoglobin (as naturally occurring chromophore present in blood vessels in the dermis). Alternatively, the superficial area may be treated with an artificial chromophore which is absorbed into the dermal layer. Such an artificial chromophore may be applied to the epidermal layer in the form of a liposome-containing topical formulation. The chromophore may then permeate through the basal layer for delivery to the dermal layer.

The laser may be arranged to scan the superficial area and/or to irradiate the dermal layer in pulses. When the laser is in pulsed mode, the pulses typically have duration of 10 μ sec to 10msec (more preferably 100 μ sec to 2msec).

It is sometimes desirable to remove part of the epidermis prior to irradiating the dermal layer according to the invention. Such epidermis removal (known as skin resurfacing) may be effected mechanically (for example by abrasion), or by means of laser radiation. When laser radiation is used for this purpose, it is typically a scanner controlled CO₂ laser source.

Preferred features of the present invention will now be described with reference to the accompanying drawings, in which:

Figure 1 is a schematic representation of the three outermost layers of mammalian skin tissue;

Figure 2 is a schematic representation of partial removal of the epidermis (skin resurfacing), which is an optional step according to the invention;

Figure 3 is a schematic illustration of the result of a prior art method of wrinkle removal, which is surgical because it involves full removal of the epidermis in a selected area and therefore exposure of the dermis and consequent second degree burning; and

Figure 4 is a schematic illustration of the result of the method according to the invention, showing that the epidermis is partially intact and the basal layer fully intact.

Referring to Figure 1, the basic skin structure of mammalian skin tissue comprises three layers, the outermost epidermis 1 which is adjacent to the basal layer 2 and then the dermis 3.

Referring to Figure 2, partial removal of an area 4 of epidermis 1 by means of CO₂ laser radiation is known as skin resurfacing. This stage represents the first step of a prior art method but is an optional step according to the invention. Both the basal layer 2 and the dermis 3 are unaffected by the laser radiation.

As shown in Figure 3, prior art method of wrinkle removal results in complete removal of an area 5 of epidermis 1 and basal layer 2 by repeated exposure to CO₂ laser radiation. Partial removal of the dermis 3 also occurs, as represented by 6, leaving the dermis exposed to air. This causes a second degree burn which is slow to heal and a risk of infection.

As shown in Figure 4, the method of wrinkle removal according to the invention results in partial removal of the epidermis 1 (this is an optional step as described in Figure 2 above) and the basal layer 2 is left intact, such that the dermis 3 is not exposed to air. Laser radiation 7 is applied to the tissue and selectively absorbed by a chromophore in the dermis 3, heating the collagen and shrinking the skin hence removing the appearance of wrinkles.

Claims

1. A method of cosmetically removing wrinkles from a superficial area of mammalian skin tissue having, in the order specified, an epidermal layer, a basal layer, and a dermal layer, which method comprises:
irradiating said dermal layer through said basal layer by means of visible or infra-red radiation, said irradiation being selected to be absorbed by a chromophore in said dermal layer such that collagen present in said dermal layer is heated, while said basal layer remains intact so as to substantially inhibit contact of said dermal layer with ambient air.
2. A method according to claim 1, wherein said irradiation is from a coherent radiation source.
3. A method according to claim 2, wherein said source comprises a ruby laser arranged to target the dermis.
4. A method according to claim 2, wherein said source comprises a dye laser of wavelength selected to target oxyhemoglobin present in blood vessels in said dermal layer.
5. A method according to claim 2, wherein said source comprises a dye laser, a ruby laser, or a semiconductor laser which scans said area of mammalian skin tissue.
6. A method according to claim 5, wherein said laser comprising said source is pulsed.
7. A method according to claim 6, wherein said pulsed laser has pulses of duration 10 μ sec to 10msec.

8. A method according to any of claims 1 to 7, in which said superficial area of mammalian skin tissue is treated with an artificial chromophore which is absorbed into said dermal layer.
9. A method according to claim 8, wherein said artificial chromophore is applied to the epidermal layer in the form of a liposome-containing topical formulation.




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 97/02322

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61N5/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	EP 0 765 673 A (ESC MEDICAL SYSTEMS) 2 April 1997 see abstract	1
P,A	EP 0 763 371 A (ESC MEDICAL SYSTEMS) 19 March 1997 see abstract	1
P,A	WO 96 33773 A (OLAVESSEN) 31 October 1996 see abstract; claim 1	1

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "A" document member of the same patent family

Date of the actual completion of the international search

14 November 1997

Date of mailing of the international search report

25/11/1997

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 851 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Taccoen, J-F

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 97/02322

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0765673 A	02-04-97	AU 6446496 A CA 2187417 A JP 9103509 A	10-04-97 30-03-97 22-04-97
EP 0763371 A	19-03-97	AU 6431096 A CA 2185196 A JP 9103507 A	20-03-97 16-03-97 22-04-97
WO 9633773 A	31-10-96	AU 5405996 A	18-11-96